

Quantum-chemical calculation of the adsorption energy of the Pb atom on the graphene

Mamatova A., Glebov A., Sadykova O.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018 Author(s). Graphene can be used as sensors to detect toxic heavy metals in peripheral blood. This requires a clear understanding of adsorption process on the graphene surface. The DFT-functional method is used to calculate adsorption energy. In the article we present the calculation results and analyse interaction of Pb atom with graphene surface. It is shown that the Pb atom in B-site position on a graphene plane has a minimum of energy.

<http://dx.doi.org/10.1063/1.5079388>

References

- [1] Brief guide to analytical methods for measuring lead in blood (World Health Organization, 2011).
- [2] A. Geim and K. Novoselov, The rise of graphene. *naturematerials*, 6: 183-191, 2007.
- [3] H. Gao, J. Zhou, M. Lu, W. Fa, and Y. Chen, *Journal of applied physics* 107, p. 114311 (2010). 10.1063/1.3437640
- [4] F. Neese, *Wiley Interdisciplinary Reviews: Computational Molecular Science* 2, 73-78 (2012). 10.1002/wcms.81
- [5] K. Nakada and A. Ishii, in *Graphene Simulation* (InTech, 2011).
- [6] K. Nakada and A. Ishii, *Migration of adatom adsorption on graphene using DFT calculation*, Vol. 151 (Elsevier, 2011), pp. 13-16.
- [7] A. D. Becke, *The Journal of chemical physics* 98, 5648-5652 (1993). 10.1063/1.464913
- [8] D. A. Pantazis and F. Neese, *Wiley Interdisciplinary Reviews: Computational Molecular Science* 4, 363-374 (2014).
- [9] F. Weigend and R. Ahlrichs, *Physical Chemistry Chemical Physics* 7, 3297-3305 (2005). 10.1039/b508541a
- [10] A. Schäfer, C. Huber, and R. Ahlrichs, *The Journal of Chemical Physics* 100, 5829-5835 (1994). 10.1063/1.467146
- [11] I. Shtepliuk, N. M. Caffrey, T. Iakimov, V. Khranovskyy, I. A. Abrikosov, and R. Yakimova, *Scientific reports* 7, p. 3934 (2017). 10.1038/s41598-017-04339-8